



Aberystwyth University

Role of live microbial feed supplements with reference to anaerobic fungi in ruminant productivity: A review

Puniya, Anil K.; Salem, Abdelfattah Z. M.; Kumar, Sanjay; Dagar, Sumit Singh; Griffith, Gareth; Puniya, Monica; Ravella, Sreenivas R.; Kumar, Nikhil; Dhewa, Tejpal; Kumar, Ravinder

Published in:

Journal of Integrative Agriculture

DOI:

[10.1016/S2095-3119\(14\)60837-6](https://doi.org/10.1016/S2095-3119(14)60837-6)

Publication date:

2015

Citation for published version (APA):

Puniya, A. K., Salem, A. Z. M., Kumar, S., Dagar, S. S., Griffith, G., Puniya, M., ... Kumar, R. (2015). Role of live microbial feed supplements with reference to anaerobic fungi in ruminant productivity: A review. *Journal of Integrative Agriculture*, 14(3), 550-560. [https://doi.org/10.1016/S2095-3119\(14\)60837-6](https://doi.org/10.1016/S2095-3119(14)60837-6)

General rights

Copyright and moral rights for the publications made accessible in the Aberystwyth Research Portal (the Institutional Repository) are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Aberystwyth Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Aberystwyth Research Portal

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

tel: +44 1970 62 2400
email: is@aber.ac.uk

Role of Live Microbial Feed Supplements with Reference to Anaerobic Fungi in Ruminant Productivity

Anil Kumar Puniya^{1*}, A Z M Salem², Sanjay Kumar^{1,3}, Sumit Singh Dagar^{1,6}, Gareth Wyn Griffith⁴, Monica Puniya¹, Sreenivas Rao Ravella⁴, Nikhil Kumar¹, Tejpal Dhewa⁵ and Ravinder Kumar¹

¹*Dairy Microbiology Division, National Dairy Research Institute, Karnal 132001, India*

²*Faculty of Veterinary Medicine and Animal Husbandry, Autonomous University of Mexico State, Mexico State C.P. 50000, Mexico*

³*Department of Clinical Studies, School of Veterinary Medicine, University of Pennsylvania, Pennsylvania 19348, USA*

⁴*Institute of Biological, Environmental and Rural Sciences, Aberystwyth University, Aberystwyth SY 23 3DD, UK*

⁵*Department of Microbiology, Bhaskaracharya College of Applied Sciences, University of Delhi, Dwarka, New Delhi 110075, India*

⁶*Microbial Science Division, Agharkar Research Institute, Pune 411004, India*

ABSTRACT

To keep the concept of a safe food supply to the consumers, animal feed industries world over are showing an increasing interest in the direct fed microbials (DFM) for improved animal performance in terms of growth or productivity. This becomes all the more essential in a situation, where a number of the residues of antibiotics and/or other growth stimulants reach in milk and meat with a number of associated potential risks for the consumers. Hence, in the absence of growth stimulants, a positive manipulation of the rumen microbial ecosystem to enhance the feedstuff utilization for improved production efficiency by ruminants has become of much interest to the researchers and entrepreneurs. A few genera of live microbes (i.e., bacteria, fungi and yeasts in different types of formulations from paste to powder) are infrequently used as DFM for the domestic ruminants. These DFM products are live microbial feed supplements containing naturally occurring microbes in the rumen. Among different DFM possibilities, anaerobic rumen fungi (ARF) based additives

*Correspondence: akpuniya@gmail.com

have been found to improve ruminant productivity consistently during feeding trials. Administration of ARF during the few trials conducted, led to the increased weight gain, milk production, and total tract digestibility of feed components in ruminants. Anaerobic fungi in the rumen display very strong cell-wall degrading cellulolytic and xylanolytic activities through rhizoid development, resulting in the physical disruption of feed structure paving the way for bacterial action. Significant improvements in the fiber digestibility were found to coincide with increases in ARF in the rumen indicating their role. Most of the researches based on DFM have indicated a positive response in nutrient digestion and methane reducing potential during *in vivo* and/ or *in vitro* supplementation of ARF as DFM. Therefore, DFM especially ARF will gain popularity but it is necessary that all the strains are thoroughly studied for their beneficial properties to have a confirmed ‘generally regarded as safe’ status for ruminants.

Keywords: anaerobic rumen fungi, bacterial DFM, direct fed microbials, probiotics, rumen

INTRODUCTION

Improved ruminant health and performance has always remained a primary objective of people associated with livestock production. Several compounds have been used to improve ruminant performance either by manipulation of the rumen environment (e.g., sodium bicarbonate) or by directly altering the composition and metabolic activities of the rumen microbes (e.g., ionophores). But, with the growing concerns towards the use of antibiotics and other growth stimulants in the ruminant feed industry, more emphasis has been given to increasing public awareness, disease prevention and use of other natural growth promoters like direct-fed microbials (DFM). DFM are the mono or mixed cultures of live microbes which when fed to the host, exert beneficial health effects by improving its gastrointestinal tract microbial balance. Aside from improving the digestibility and performance of the ruminants, DFM detoxify toxic compounds to modulate immune system and maintain gut peristalsis and intestinal mucosal integrity (Chaucheyras-Durand and Duran 2010, Sandri *et al.* 2014). The term DFM is different from “Probiotic” in a sense that it is only restricted to the use of “live, naturally occurring microbes” (Yoon and Stern 1995; Krehbiel *et al.* 2003; Kenney 2013). For domestic ruminants like cattle and buffaloes, yeasts and aerobic fungi have been successfully used to increase growth rate and production efficiency. But, now a day’s use of anaerobic fungi is emphasized because of its ability to produce wide array of enzymes that can even degrade the lignified walls of plant-cells. Many factors like infections, improper food, environmental conditions and ingestion of antibiotics have been described that result in imbalance

of intestinal microflora of ruminants. For many years, studies related to supplementation of microbial feed additive in the diet for the improvement of health are under progress. Now days, there are growing evidences that DFM may be useful in managing conditions like irritable bowel syndrome, lactose intolerance, chronic liver disease, pancreatitis and even certain forms of cancers. The mechanisms suggested for the action for DFM include colonization of the lower intestine, thereby limiting the growth of any potential pathogens through ‘competitive exclusion’ or inhibit pathogens by lowering the pH of the intestinal lumen and by producing anti-microbial proteins (bacteriocins).

This paper will cover a number of aspects related to the type of DFM, their mode of action, environmental protection using DFM, their benefits when fed to the host etc.

BACTERIAL DFM

There are many DFM based on bacteria that are commercially available for use in ruminant diets with more specific applications. Most of the DFM bacteria are lactic acid bacteria with lactobacilli being the most dominant microflora, followed by the bifidobacteria, enterococci and bacilli. Among lactobacilli, *Lactobacillus acidophilus* is the most commonly used in DFM. Most bacterial-based DFM are probably beneficial because they have effects in the lower gut and not in the rumen. For example, *L. acidophilus* produces lactic acid, which may lower the pH in small intestines, and inhibit the growth of pathogenic microbes. Early research with DFM was focused on ruminants which are either stressed or having immature microbial ecosystems in their guts (Vandevoorde *et al.* 1991) like milk fed young calves, calves being weaned or cattle being shipped (Jenny *et al.* 1991).

Modes of action

In ruminants, mode of action of feeding bacterial DFM is variable, which emphasizes the need for greater understanding of underlying mechanisms. Research conducted to determine the potential mode of action of bacterial DFM has most often used the rodent models. Bacterial DFM have been reported to modify the balance of intestinal microbes, adhere to intestinal mucosa and prevent pathogen adherence or activation, influence gut permeability, and modulate immune function are discussed below.

Competitive attachment Early research (Jones and Rutter 1972) suggested that attachment to the intestinal wall was important for pathogenic strains of *E. coli* to induce diarrhea. It is believed that the attachment support proliferation and reduce peristaltic removal of organisms. Bacterial DFM could compete with pathogens for the sites of adherence on the intestinal surface and thus can

facilitate their removal (Wisener *et al.* 2014). Adhesion is thought to be mediated either nonspecifically by physicochemical factors, or specifically by adhesive bacterial surface molecules and epithelial receptor molecules (Holzapfel *et al.* 1998).

Antibacterial effect Many species of lactobacilli have demonstrated inhibitory activity against pathogens. *L. acidophilus* has been shown to be antagonistic toward entero-pathogenic *E. coli*, *Salmonella typhimurium*, *Staphylococcus aureus* and *Clostridium perfringens* (Gilliland and Speck 1977). Mann *et al.* (1980) showed that the strain of *E. coli*, which causes illness and death when it is the sole microbial species in young lambs, could be tolerated in the presence of lactobacilli. Hydrogen peroxide produced by lactobacilli appears to be partially responsible for the antagonistic interaction (Gilliland and Speck 1977). Different reports suggest that antimicrobial proteins and/or bacteriocins either mediate or facilitate antagonism by *L. acidophilus* (Gilliland and Speck 1977; Barefoot and Klaenhammer 1983). However, because of the presence of proteolytic enzymes, their importance might be limited. In addition, Walsh *et al.* (2012) suggested that DFM should not be considered as viable alternatives to in-feed antibiotics in a pathogen challenge situations.

Immune Response Bacterial DFM have been shown to affect the innate, humoral and cellular arms of the immune system. Oral administration of lactobacilli generally result in an augmentation of innate immune responses (i.e., enhanced phagocytosis and natural killer cell activity), as well as an elevate production of immunoglobulin (IgA) and a decrease IgE production in animals (Erickson and Hubbard 2000; Isolauri *et al.* 2001). However, influence of DFM on cytokine production and T and B cell responses show mixed results depending on the strain, dose and duration of feeding DFM, as well as the type of tissues and cells analyzed. Furthermore, some species of probiotics appear to be capable of altering the immunomodulatory effects exerted by other species. For instance, *L.reuteri* DSM12246 was shown to potentially suppress *Lactobacillus. casei* induced production of IL-6, IL- 12, and TNF- α in dendritic cells (Christensen *et al.* 2002), suggesting that the composition of bacterial DFM administered should be considered. Qiu *et al.* (2012) indicated that supplementation with the DFM also regulate in energy re-partitioning to the immune system and an increase in antibody production independent of changes in whole body metabolism or growth performance. Therefore, bacterial DFM also show promise as immune modulators, although, more research is needed to determine the underlying mechanisms.

Effect on ruminant performance

Pre-ruminant calves Generally, the importance of feeding DFM to neonatal livestock has been to establish and maintain normal intestinal microbes rather than as a production stimulant. In the

neonate, the microbial population of the gastrointestinal tract (GIT) is in transition and extremely sensitive. Abrupt environmental or dietary changes may cause shifts in the microbial population of the GIT which often leads to an increased incidence of diarrhea in calves (Sadine 1979). In terms of ruminant production systems, the efficacy of bacterial DFM has been studied most extensively in the neonatal dairy calf. Bacterial DFM, such as species of *Lactobacillus*, *Enterococcus*, *Streptococcus*, and *Bifidobacterium* have been studied in young calves and the data have been reviewed. For dairy calves, rapid adaptation to solid feed by accelerating the establishment of rumen and intestinal microbes and avoiding the establishment of entero-pathogens, which often results in diarrhea, is the primary goal. Feeding calves with viable cultures of species of *Lactobacillus* and *Streptococcus* has been reported to decrease the incidence of diarrhea (Ewaschuk *et al.* 2004; Hossaini *et al.* 2010; Riddell *et al.* 2010). In addition, some studies have indicated that DFM in the diet improves weight gain, feed efficiency and feed intake (Timmerman *et al.* 2005; Adams *et al.* 2008). In an experiment by Hossaini *et al.* (2010), calves fed DFM containing *L. acidophilus*, *L. casei*, *Bacillus. thermophilus*, *Enterococcus. faecium* confirmed the beneficial effect of it. The decreased incidence of diarrhea might be associated with a consistently increased shedding of *Lactobacillus* (Gilliland *et al.* 1980; Jenny *et al.* 1991; Abu-Tarboush *et al.* 1996) and an inconsistent decreased shedding of coliforms (Bruce *et al.* 1979) in feces in response to supplements of *Lactobacillus*.

Performance response is likely not important early in the pre-ruminant's life when enteric disease is most prevalent. Improved health and reduction in the incidence or severity of diarrhea, though difficult to measure for statistical analysis, is most likely a more important response. As suggested by Newman and Jacques (1995), more experiments that include detailed information about the microbial supplement, and fecal culture data from scouring experimental animals are needed to determine the usefulness of microbial supplements in neonatal calves.

Lactating Ruminants Modern day intensive production systems, especially with high producing dairy cows and buffaloes involve the feeding of high levels of concentrate in order to meet the metabolic demand for high milk yield. Feeding high levels of concentrate often lead to metabolic dysfunction and eventually rumen acidosis; especially under conditions of poor methods of feeding and/or composition of diets. The goal of the nutritionist, when implementing high concentrate feeding is to maximize performance and efficiency, while keeping digestive disturbances such as the rumen acidosis within acceptable limits through good nutritional management. Theoretically, a number of approaches can be followed to control the incidences of the rumen acidosis. One approach is to inhibit the growth of lactic acid producing bacteria such as *Streptococcus bovis* and

Lactobacillus species through the use of feed supplements such as ionophores (Callaway and Martin 1997). Another approach is to use DFM such as *Megasphaera elsdenii*, a lactic acid utilizer, to regulate lactic acid levels in the rumen. Experimentally, there have been several bacteria that have potential as DFM for ruminants but have not been commercialized for different reasons. For example, *M. elsdenii* is the major lactate-utilizing organism in the rumen of adapted cattle fed high grain diets. However, when cattle are abruptly shifted from a high-forage to high concentrate diet, the numbers of *M. elsdenii* are often insufficient to prevent lactic acidosis. Similarly, *E. faecium* and yeast used were of limited value for feedlot cattle already adapted to high-grain diets (Beauchemin *et al.* 2003). Erasmus (1992) and Aikman *et al.* (2008) observed an increase in milk production for a high producing group of cows when *M. elsdenii* NCIMB 41125 was dosed compared to the control animals. Similar results were obtained in second lactating cows (Hagg and Henning 2007), where *M. elsdenii* NCIMB 41125 were dosed after calving.

Gomez-Basauri *et al.* (2001) reported 0.73 kg/d more milk with 0.42 kg less DM consumption, when cows were fed with lactic acid bacteria (*L. acidophilus*, *L. casei*, *E. faecium*; total lactic bacteria=10⁹ cfu g⁻¹) and mannan-oligosaccharide, compared to control. Furthermore, milk yields continued to increase over time for DFM- and mannan-oligosaccharide-fed cows, whereas control cows maintained constant milk yields. On similar lines, Boyd *et al.* (2011) reported that the addition of a direct-fed microbial (*L. acidophilus* NP51 and *Proponibacterium freudenreichii* NP24) and dietary glycerol may improve yield and digestibility for cows subject to heat stress. However, strain difference (*L. acidophilus* LA747 and *Proponibacterium freudenreichii* PF24) may not affect the performance, diet digestibility and rumen characteristics (Raeth-Knight *et al.* 2007).

Other experiments conducted with combinations of fungal cultures and lactic acid bacteria (Komari *et al.* 1999; Block *et al.* 2000) has shown higher milk yields when lactating cows were fed with *Saccharomyces cerevisiae* in combination with *L. acidophilus* and/or *Lactobacillus plantarum*/*E. faecium*. Propionibacteria, which convert lactic acid and glucose to acetic and propionic acid, may also be beneficial if inoculated into the rumen, because higher concentrations of rumen propionate represents the energy status of the animal. These bacteria are naturally present in high numbers in the rumen of animals fed forage and medium concentrate diets. Their supplementation as DFM increased milk fat percentage and milk yield as well as improved health of prepartum and postpartum cows (Noeck *et al.* 2006; Oetzel *et al.* 2007).

YEAST AND FUNGAL DFM

In adult ruminants, fungal DFM have mostly been selected to target the rumen compartment, which is the main site for feed digestion. The fungal feed additives and supplements have been shown to affect the rumen fermentation patterns.

Mode of action

Several reasons for improvements in rumen fermentation from feeding fungal DFM have been suggested. First, DFM exerts beneficial changes in activity and numbers of the rumen microbes. For example, the total rumen anaerobes and cellulolytic bacteria increase with fungal extracts. Beharka *et al.* (1991) reported that young calves fed *Aspergillus oryzae* fermentation extract were weaned one week earlier than untreated calves and that supplementation increased the rumen bacteria and VFA concentrations. *Aspergillus* fermentation extracts (Chang *et al.* 1999) and yeast cultures (Chaucheryas *et al.* 1995) have also been shown to stimulate the rumen fungi directly, which improved fiber digestion. Feeding *S. cerevisiae* increased the rumen protozoa and increased NDF digestion in steers fed straw-based diets (Plata *et al.* 1994). Yeasts have also been shown to stimulate acetogenic bacteria in the presence of methanogens (Chaucheryas *et al.* 1995), which might result in more efficient rumen fermentation.

Second, fungal DFM may also prevent the accumulation of excess lactic acid in the rumen when cattle are fed diets containing highly fermentable carbohydrates. Specifically, extracts of *A. oryzae* stimulated the uptake of lactic acid by the rumen lactate-utilizers *Selenomonas ruminantium* (Nisbet and Martin 1991) and *M. elsdenii* (Waldrup and Martin 1993) possibly by providing a source of malic acid. Increased metabolism of lactic acid should theoretically raise rumen pH and this may be one reason why DFM increased the rumen cellulolytic bacteria and improved fiber digestion (Arambel *et al.* 1987). Chaucheyras *et al.* (1995) reported that *S. cerevisiae* was able to prevent the accumulation of lactic acid production by competing with *S. bovis* for glucose and by stimulating the uptake of lactic acid by *M. elsdenii*, perhaps by supplying amino acids and vitamins. In contrast, added yeasts were unable to prevent acute episodes of lactic acidosis when fermentations were challenged with a diet rich in fermentable carbohydrates (Aslan *et al.* 1995). Yeast may improve rumen fermentation because they are able to scavenge excess oxygen (Newbold *et al.* 1996), creating a more optimal environment for the rumen anaerobic bacteria. *Aspergillus* extracts may improve fiber digestion because they contain esterase enzymes (Varel *et al.* 1993).

Anaerobic rumen fungi (ARF) have also been supplemented as fungal DFM to ruminant for better utilization of fibrous feeds in terms of increased feed intake, body weight gain, enhanced milk production, and thus improved ruminant productivity (Dey *et al.* 2004; Thareja *et al.* 2006). ARF

are the normal inhabitants of the rumen ecosystem. The fungi colonize the fibrous plant fragments in the rumen and penetrate plant tissues making more room for bacterial attack and thus increase the area susceptible to enzymatic attack (Dagar *et al.* 2011). The enzymes produced by ARF and their functions are shown in table 1. These properties of ARF are suggestive of manipulation of fungal numbers for better utilization of fibrous feeds.

Effect on ruminant performance

There have been numerous studies reporting positive effects of *S. cerevisiae* and *A. oryzae* on intake and milk production of lactating cows. Supplementing diets with *S. cerevisiae* was shown to increase total dry matter intake, total volatile fatty acids (VFA) and propionic acid production, besides higher propionate concentration and decreased acetate to propionate ratio were determined in some experiments (Schingoethe *et al.* 2004; Ondarza *et al.* 2010; Cakiroglu *et al.* 2010). Higher VFA, especially propionic acid are important in terms of enhanced lactose production, milk volume and overall energy balance (Miller-Webster *et al.* 2009). Erasmus *et al.* (1992) suggested that supplementation of *S. cerevisiae* tended to increase microbial protein synthesis in dairy cows and significantly altered the amino acid profile of the duodenal digesta. Wohlt *et al.* (1991) suggested that supplementing yeast culture before parturition and extending through peak lactation was necessary to evaluate the effect on lactating cows. Some field reports indicate increased dry matter intake (DMI) and milk production when yeast was fed during periods of heat stress, possibly reflecting the role in aiding appetite during time of stress (Huber, 1998). In beef cattle the addition of *S. cerevisiae* lead to an increase of live weight by 7.5% depending on the type of diet tested. Improvement can reach 13% in feedlot conditions, with diets rich in starch and sugars. Wallace and Newbold (1993) reported that responses recorded in trials in beef cattle tended to be higher with corn silage rather than with grass silage. In dairy cows, an improvement by around 4% of the milk yield, often associated with increased feed intake was generally reported and response was greater in early as opposed to mid or late lactation (Ali-Haimoud-Lekhal *et al.* 1999). *A. oryzae* in diets of lactating cows increased milk production, feed efficiency and tolerance to heat stress in some (Gomez-Alarcon *et al.* 1990) but not all (Higginbotham *et al.* 1993; Yu *et al.* 1997) studies.

Among microbial additives, there are evidences of definite positive relationship between ARF in the rumen and the increased voluntary intake of low digestible fibrous feeds (McAllister *et al.* 1994; Ha *et al.* 1994; Dey *et al.* 2004; Saxena *et al.* 2010). The ARF have been isolated from animals of different parts of the world providing evidence to suggest that they may have an important role in the digestion of fibrous materials in the rumen (Trinci *et al.* 1994; Tripathi *et al.* 2007b; Dagar *et al.*

2011; Ishtiyak *et al.* 2013) through substantial colonization of plant material (Edwards *et al.* 2008). Different fungal species improved digestibility of dry matter and cell wall constituents of cereal straws (Manikumar *et al.* 2004) as well as sugarcane bagasse (Shelke *et al.* 2009) in the *in vitro* system. Incorporation of fungus increased growth rate, rumen fermentation, nutrient digestibility and nitrogen retention in sheep (Ha *et al.* 1994), crossbred calves (Dey *et al.* 2004), and buffalo calves (Sehgal *et al.* 2008). Tripathi *et al.* (2007a, b) found that administration of *Piromyces* sp. increased the growth rate, feed efficiency and nutritive value of wheat straw based ration in buffalo calves.

Experiments, where ARF were either absent or eliminated have provided a deep insight into the contribution of fungi to fibre digestion, feed intake, rumen fermentation and overall metabolism. Ford *et al.* (1987) showed a decrease in voluntary feed intake of sheep to 49% in groups where ARF were eliminated. Removal of ARF from the rumen of sheep reduced the voluntary intake of poor quality feed to about 70% (Gordon and Phillips 1993). The addition of fungal culture *Neocallimastix* sp. R1 increased the forage intake by 35% in early weaned calves (Theodorou *et al.* 1990). In fungi free rumen of sheep, the dosing of *Neocallimastix* sp. SL1 increased the intake of straw based diet to 40% (Gordon and Phillips 1993). The elimination of ARF significantly reduced the degradation of dry matter, neutral detergent fiber, acid detergent fiber, and the activity of CMCase in sheep rumen (Gao *et al.* 2013).

An increased feed digestibility was documented, when different strains of *Neocallimastix* were dosed into the rumen of fungus free sheep (Elliott *et al.* 1987). Paul *et al.* (2004) studied the effect of *Piromyces* sp. FNG5 on *in vivo* rumen fermentation and digestion of nutrients in buffaloes. They found an increase in total tract DMD, organic matter, neutral detergent fibre and acid detergent fibre digestibility. An increase in VFAs and enzymatic activities (carboxymethylcellulase (CMCase), xylanase, microcrystalline cellulase, acetyl esterase, feruloyl esterase and protease) was also noticed. In addition, *Piromyces* sp. FNG5 was also found to tolerate tannic acid concentration up to 20 g/L (Paul *et al.* 2006), suggesting its possible application in improving fibre digestion of tannin-containing feeds. The administration of ARF into the rumen of goat increased the DMD, concentrations of ammonia, total VFA and CMCase activity. On the other hand, their elimination from sheep and goat resulted in a decreased digestibility of straw based dry matter. In absence of ARF, the concentrations of acetate, butyrate and total VFA decreased significantly in the rumen of sheep (Gao *et al.* 2008). Sehgal *et al.* (2008) studied the influence of *Neocallimastix* sp. GR1 on growth, rumen fermentation and nutrient digestion in female buffalo calves and found a considerable increase in daily weight gain and better feed efficiency of total mixed ration compared

to control calves. Tripathi *et al.* (2007b) found that the DMD was highest in group fed with *Piromyces* sp. WNG-12 than *Orpinomyces* sp. C-14 fed group. A similar pattern of increased digestibility of crude protein, cell-wall contents and average body weight gain was also observed in treatment groups. The same cultures were used to study the digestibility of wheat straw: concentrate (50:50) based diet, effect on rumen fermentation and milk production in lactating buffalo (Saxena *et al.* 2010). An increase in milk production was recorded in the fungus fed groups. There was also an increase of 6% fat corrected milk yield/ animal/ day in treatment groups. A similar pattern of increase in DMD, crude protein, neutral detergent fibre, acid detergent fibre, cellulose and digestible energy were observed in fungus fed groups, extending the possibility of their use as DFM in lactating buffaloes for obtaining higher milk production, even on poor quality feed.

ENVIRONMENTAL PROTECTION USING DFM

Methane produced from enteric fermentation leads to loss of 6 to 15% of gross intake energy of ruminant's energy. Besides, methane is the second most potent green house gas, lead to the global warming and poses threats to the environment (Kumar *et al.* 2009, 2013a, b, 2014). Thus, the consequences of methanogenesis in the rumen is not only associated with low ruminant efficiency but also have a negative impact on the sustainability of their production. Since, the enteric fermentation emission is one of the major sources of methane; therefore, experiments were conducted using antibiotics and other chemicals for mitigating methane emissions. However, appearance of antibiotic-resistant bacteria restricts its convenient use. Moreover, the antibiotics excreted to manures without being absorbed have been scattered on the environment (Mwenya *et al.* 2006). The alternative to antibiotics is the use of DFM that include lactic acid bacteria and yeasts as they are also found to reduce methane emission (Kalmakoff *et al.* 1996; Teather and Forster 1998; Klieve and Hegarty 1999) and acetate: propionate ratio (Martin and Nisbet 1992; Gamo *et al.* 2002; Lila *et al.* 2004). Hydrogen, which is released in the rumen during fibre degradation by cellulolytic microbes like bacteria and ARF, is rapidly utilized by methanogens for its conversion to methane. On the other hand, acetogenic bacteria are also able to utilize hydrogen for acetate production; but their number is less in the rumen of adults. Therefore, the acetogenic bacteria could be potentially used to compete with methanogens for hydrogen utilization; thereby also preventing the energy loss occurring as a result of methane production. Chaucheyras *et al.* (1995) studied the effect of a live strain of *S. cerevisiae* on hydrogen utilization and acetate and methane production by an acetogen and a methanogen. They concluded that the addition of yeast cells enhanced the acetogenesis of the acetogenic strain by more than fivefold, while in absence of yeasts, hydrogen was principally used

for methane synthesis. Therefore, the use of yeasts as ruminant feed additives could help reducing methane, increasing the rumen metabolism and hence, promoting ruminant performance and health. Lopez *et al.* (1999) also found that acetogens depress methane production when added to the rumen fluid *in vitro* and suggested that even if a stable population of acetogens could not be established in the rumen, it might be possible to achieve the same metabolic activity using the acetogens as a daily fed feed additive. In addition, methane oxidisers can also be used as DFM. The oxidation reaction competes with the production of methane, which is a strictly anaerobic process. Methane oxidisers from gut and non-gut sources could be screened for their activity in the rumen fluid *in vitro* and then selected methane oxidisers could be introduced into the rumen on a daily basis.

PRACTICAL APPLICATIONS OF DFM

There are varieties of DFMs such as powder, paste, gel, and capsules available commercially. These different forms may be mixed in feed, top-dressed, given as a paste, or mixed into the drinking water or milk replacer. However, their use must be managed effectively as viability of organism can be largely affected on interactions with chlorine, water, temperature, minerals, flow rate, and antibiotics. Bacterial DFM pastes are formulated with vegetable oil and inert gelling ingredients. Non-hydroscopic whey is generally used as a carrier for bacteria based DFM. Fungal DFM products are formulated with grain by-products as carriers. Some DFM are developed for one-time dosing while others are developed for feeding on a daily basis. Most DFMs contain live bacteria; however, some contain only bacterial or fungal extracts or fermentation by-products. The best response can be observed during the following situations: (a) when a newborn animal acquire beneficial bacteria from environment, (b) during weaning or dietary changes, (c) periods of stress i.e. shipping, vaccination, and other situations , and (d) antibiotic therapy. The stability of DFMs is crucial because the microbes must be delivered live to the animal to be effective. For this, most DFMs require storage in a cool and dry area, away from heat, direct sunlight, and high levels of humidity. They must not only survive during processing and storage but also in the gut environment. The metabolites present in culture extracts have been suggested to be the “active” ingredients.

CONCLUSION AND FUTURISTIC APPROACHES

In light of international regulations and consumer demands to withdraw the growth-enhancing antibiotics and limiting the use of treatment related antibiotics, the DFM offer an option. For ruminants, ARF as DFM have been used successfully for improving the rumen and gastro-intestinal health, enhancing milk production, feed efficiency and daily gain in animals. On the other hand,

methanogenesis, which accounts for significant loss of ruminant's energy and increased greenhouse gases in environment, is also a major concern in present scenario. Therefore, the use of DFM for improving production efficiency without compromising animal health and environmental sustainability is most advocated.

Acknowledgements

We thankfully acknowledge the DBT-CREST fellowship 2011-12 to Anil K Puniya that greatly helped in developing the manuscript in collaboration with our overseas expert GWG in UK and elsewhere. We also acknowledge the financial support provided under the Network Project of ICAR on 'VTCC' to carry the research further in this direction.

References

- Abu-Tarboush H M, Al-Saiady M Y, Keir El-Din A H. 1996. Evaluation of diet containing lactobacilli on performance, fecal coliform, and lactobacilli of young dairy calves. *Animal Feed Science and Technology*, **57**, 39–49.
- Adams M C, Luo J, Rayward D, King S, Gibson R, Moghaddam G H. 2008. Selection of a novel direct-fed microbial to enhance weight gain in intensively reared calves. *Animal Feed Science and Technology*, **145**, 41–52.
- Aikman P C, Henning P H, Jones A K, Potteron S, Siviter J, Carter S, Hill S, Kirton P, Szoka R. 2008. Effect of administration of *Megasphaera elsdenii* NCIMB 41125 lactate utilising bacteria in early lactation on the production, health and rumen environment of highly productive dairy cows fed a high concentrate diet. *Animal Nutrition Internal Report*.
- Ali-Haimoud-Lekhal D, Lescoat P, Bayourthe C, Moncoulon R. 1999. Effect of *Saccharomyces cerevisiae* and *Aspergillus oryzae* on milk yield and composition in dairy cows: A review. *Rencontres Recherche Ruminants*, **6**, 157.
- Arambel M J, Weidmeier R D, Walters J L. 1987. Influence of donor animal adaptation to added yeast culture and/or *Aspergillus oryzae* fermentation extract on *in vitro* rumen fermentation. *Nutritional Reports International*, **35**, 433–437.
- Aslan V S, Thamsborg M, Jorgensen R J, Basse A. 1995. Induced acute ruminal acidosis in goats treated with yeast (*Saccharomyces cerevisiae*) and bicarbonate. *Acta Veterinaria Scandinavica*, **36**, 65–68.
- Atanasova-Pancevska N, Kungulovski D. 2008. Comparison of morphological and enzyme characteristics of anaerobic fungi isolated from *Cervus dama*. *Central European Journal of Biology*, **3**(1), 69–74.
- Atsushi K, Azuma J-I, Koshijima T. 1984. Lignin-carbohydrate complexes and phenolic acids in bagasse. *Holzforschung*, **38**(3), 141–149.

- 393 Barefoot S F, Klaenhammer T R. 1983. Detection and activity of lactacin B, a bacteriocin produced by
394 *Lactobacillus acidophilus*. *Applied and Environmental Microbiology*, **45**, 1808–1815.
- 395 Beauchemin K A, Yang W Z, Morgavi D P, Ghorbani, G R, Kautz W, Leedle J A Z. 2003. Effects of bacterial
396 direct-fed microbials and yeast on site and extent of digestion, blood chemistry, and subclinical ruminal
397 acidosis in feedlot cattle. *Journal of Animal Science*, **81**, 1628-1640.
- 398 Beharka A A, Nagaraja T G, Morrill J L. 1991. Performance and ruminal development of young calves fed diets
399 with *Aspergillus oryzae* fermentation extracts. *Journal of Dairy Science*, **74**, 4326-4336.
- 400 Block E, Nocek J E, Kautz W P, Leedle J A Z. 2000. Direct fed microbial and anionic salt supplementation to
401 dairy cows fed 21 days pre- to 70 days postpartum. *Journal of Animal Science*, **78**, 304.
- 402 Blum DL, Li X-L, Chen H, Ljungdahl LG. 1999. Characterization of an acetyl xylan esterase from the anaerobic
403 fungus *Orpinomyces* sp. Strain PC-2. *Applied and Environmental Microbiology*, **65(9)**, 3990-3995.
- 404 Boyd J, West J. W, Bernard J K. 2011. Effects of the addition of direct-fed microbials and glycerol to the diet of
405 lactating dairy cows on milk yield and apparent efficiency of yield. *Journal of Dairy Science*, **94**, 4616-4622.
- 406 Breton A, Gaillard-Martinie B, Gerbi C, Gomez de Segura B, Durand R, Kherratia B. 1995. Location by
407 fluorescence microscopy of glycosidases and a xylanase in the anaerobic gut fungi *Caecomyces communis*,
408 *Neocallimastix frontalis*, and *Piromyces rhizinflata*. *Current Microbiology*, **31(4)**, 224-227.
- 409 Bruce B B, Gilliland S E, Bush L J, Staley T E. 1979. Influence of feeding cells of *Lactobacillus acidophilus* on
410 the fecal flora of young dairy calves. *Oklahoma Animal Science Research Report*, Stillwater, OK, p. 207
- 411 Cakiroglu D, Meral Y, Pekmezci D, Akdag F. 2010. Effects of live yeast culture (*Saccharomyces cerevisiae*) on
412 milk production and blood lipid levels of cows in early lactation. *Journal of Animal and Veterinary Advance*,
413 **9**, 1370-1374.
- 414 Callaway T R, Martin S A. 1997. Effects of cellobiose and monensin on *in vitro* fermentation of organic acids by
415 mixed ruminal bacteria. *Journal of Dairy Science*, **80**, 1126-1135.
- 416 Chang J S, Harper E M, Calza R E. 1999. Fermentation extract effects on the morphology and metabolism of the
417 rumen fungus *Neocallimastix frontalis* EB188. *Journal of Applied Microbiology*, **86**, 389-398.
- 418 Chaucheyras F, Fonty G, Bertin G, Gouet P. 1995. *In vitro* H₂ utilization by a ruminal acetogenic bacterium
419 cultivated alone or in association with archaea methanogen is stimulated by a probiotic strain of
420 *Sacharomyces cerevisiae*. *Applied and Environmental Microbiology*, **61**, 3466-1995.
- 421 Chaucheyras-Durand F, Duran H. 2010. Probiotics in animal nutrition and health. *Beneficial Microbes*, **1**, 3-9.
- 422 Christensen H R, Frokiaer H, Pestka J J. 2002. Lactobacilli differentially modulate expression of cytokines and
423 maturation surface markers in murine dendritic cells. *Journal of Immunology*, **168**, 171–178.
- 424 Comlekcioglu O, Ozkose E, Tutus A, Akyol I, Ekinci M S. 2010. Cloning and characterization of cellulase and
425 xylanase coding genes from anaerobic fungus *Neocallimastix* sp. GMLF1. *International Journal of*
426 *Agriculture and Biology*, **12(5)**, 691-696.
- 427 Coughlan M P, Hazlewood G P. 1993. beta-1,4-D-xylan-degrading enzyme systems: biochemistry, molecular
428 biology and applications. *Biotechnology and Applied Biochemistry*, **17(3)**, 259-289.

- 429 Dagar S S, Kumar S, Mudgil P, Singh R, Puniya A K. 2011. Use of D1/D2 domain of large subunit rDNA as a
430 taxonomic marker and differentiation of *Orpinomyces* spp. using PCR-RFLP. *Applied and Environmental*
431 *Microbiology*, **77**, 6722-6725.
- 432 Dey A, Sehgal J P, Puniya A K, Singh K. 2004. Influence of an anaerobic fungal culture (*Orpinomyces* sp.)
433 administration on growth rate ruminal fermentation and nutrient digestion in calves. *Asian-Australasian*
434 *Journal of Animal Science*, **17**, 820-824.
- 435 Edwards J E, Kingston-Smith A H, Jimenez H R, Huws S A, Skot K P, Griffith G W, McEwan N R., Theodorou
436 M K. 2008. Dynamics of initial colonization of nonconserved perennial ryegrass by anaerobic fungi in the
437 bovine rumen. *FEMS Microbiology Ecology*, **66**, 537-545.
- 438 Elliott R, Ash A J, Calderon-Cortes F, Norton B W, Bauchop T. 1987. The influence of anaerobic fungi on rumen
439 volatile fatty acid concentrations *in vivo*. *The Journal of Agricultural Science*, **109**, 13-17.
- 440 Erasmus L J, Botha P M Kistner A. 1992. Effect of yeast culture supplement on production, rumen fermentation
441 and duodenal nitrogen flow in dairy cows. *Journal of Dairy Science*, **75**, 3056-3065
- 442 Erickson K L, Hubbard N E. 2000. Probiotic immunomodulation in health and disease. *American Society of*
443 *Nutritional Science*, **130**, 403S-490S.
- 444 Ewaschuk J B, Naylor J M, Chirino-Trejo M, Zello G A. 2004. *Lactobacillus rhamnosus* strain GG is a potential
445 probiotic for calves. *Canadian Journal of Veterinary Research*, **68**, 249-53.
- 446 Ford C W, Elliott R, Maynard P J. 1987. The effect of chlorite delignification on digestibility of some grass forage
447 and on intake and rumen microbial activity in sheep fed barley straw. *Journal of Agricultural Science*
448 *Cambridge*, **108**, 129-136.
- 449 Gamo Y, Mii M, Zhou X G, Sar C, Santoso B, Arai I, Kimura K, Takahashi J. 2002. Effects of lactic acid bacteria,
450 yeasts and galactooligosaccharide supplementation on *in vitro* rumen methane production. In: Takahashi J,
451 Young B A (Eds.), *Greenhouse Gases and Animal Agriculture*. Amsterdam, The Netherlands: Elsevier
452 Science BV, pp. 201-204.
- 453 Gao A, Hou X, Yang J, Fu Q. 2008. Effects of elimination of anaerobic fungi in sheep on the microbes and
454 ruminal fermentation. *Journal of Anhui Agricultural University*, **35**, 499-506.
- 455 Gao A, Wang H, Yang J, Shi C. 2013. The effects of elimination of fungi on microbial population and fiber
456 degradation in sheep rumen. *Applied Mechanics and Materials*, **295-298**, 224-231.
- 457 Gilliland S E, Bruce B B, Bush L J Staley T E. 1980. Comparison of two strains of *Lactobacillus acidophilus* as
458 dietary adjuncts for young calves. *Journal of Dairy Science*, **63**, 964-972.
- 459 Gilliland S E, Speck M L. 1977. Antagonistic action of *Lactobacillus acidophilus* toward intestinal and food
460 borne pathogens in associative cultures. *Journal of Food Protection*, **40**, 820-823.
- 461 Gomez-Alarcon R A, Dudas C, Huber J T. 1990. Influence of *Aspergillus oryzae* on rumen and total tract
462 digestion of dietary components. *Journal of Dairy Science*, **73**, 703-710.

- 463 Gomez-Basauri J, de Ondarza M B, Siciliano-Jones J. 2001. Intake and milk production of dairy cows fed lactic
464 acid bacteria and mannanoligosaccharide. *Journal of Dairy Science*, **84** (Suppl. 1), 283.
- 465 Gordon G L R, Phillips M W. 1998. The role of anaerobic gut fungi in ruminants. *Nutrition Research Review*
466 **11**(1), 133-168.
- 467 Gordon G L R, Phillips M W. 1993. Removal of anaerobic fungi from the rumen of sheep by chemical treatment
468 and the effect on feed consumption and *in vivo* fibre digestion. *Letters in Applied Microbiology*, **17**(5), 220-
469 223.
- 470 Ha J K, Lee S S, Kim C H, Choi Y J, Min H K. 1994. Effect of fungal inoculation on ruminal fermentation
471 characteristics enzyme activities and nutrient-digestion in sheep. *Proceedings of Society of Nutritional*
472 *Physiology*, **3**, 197.
- 473 Hagg F M, Henning P H. 2007. Evaluation of supplementation with *Megasphaera elsdenii* NCIMB 41125, a
474 lactate utilizing rumen microorganism, on performance in Holstein dairy cows. *KK Animal Nutrition Internal*
475 *Report*.
- 476 Higginbotham G E, Bath D L, Butler L J. 1993. Effect of feeding *Aspergillus oryzae* extract on milk production
477 and related responses in a commercial dairy herd. *Journal of Dairy Science*, **76**, 1484–1489.
- 478 Holzapfel W H, Haberer P, Snel J, Schillinger U, Huisin't Veld J H J. 1998. Overview of gut flora and probiotics.
479 *International Journal of Food Microbiology*, **41**, 85–101.
- 480 Hossaini S M R, Bojarpour M, Mamouei M, Asadian A, Fayazi J. 2010. Effects of probiotics and antibiotic
481 supplementation in daily milk intake of newborn calves on feed intake body weight gain, fecal scores and
482 health condition. *Journal of Animal and Veterinary Advance*, **9**, 872-875.
- 483 Huber J T. 1998. Yeast products help cattle handle heat. *Hoard's Dairyman*, 143:367.
- 484 Isolauri E, Sutas Y, Kankaanpää Y P, Arvilommi H, Salminen S. 2001. Probiotics: Effects on immunity.
485 *American Journal of Clinical Nutrition*, **73** (Suppl. 2), 444S–450S.
- 486 Jenny B F, Vandijk H J, Collins J A, 1991. Performance and fecal flora of calves fed a *Bacillus subtilis*
487 concentrate. *Journal of Dairy Science*, **74**, 1968–1973.
- 488 Jones G W, Rutter J M. 1972. Role of K88 antigen in the pathogenesis of neonatal diarrhoea caused by
489 *Escherichia coli* in piglets. *Infection and Immunity*, **6**, 918–927.
- 490 Kalmakoff M L, Barlett F, Teather R M. 1996. Are ruminal bacteria armed with bacteriocin? *Journal of Dairy*
491 *Science*, **79**, 2297–2306.
- 492 Kenney N. 2013. Impact of direct-fed microbials on nutrient utilization in beef cattle. Thesis and Dissertations-
493 Animal and Food Sciences. University of Kentucky, UKnowledge, UK.
- 494 Klieve A V, Hegarty R S. 1999. Opportunities for biological control of methanogenesis. *Australian Journal of*
495 *Agricultural Research*, **50**, 1315–1319.
- 496 Komari R K, Reddy Y K L, Suresh J, Raj D N. 1999. Effect of feeding yeast culture (*Saccharomyces cerevisiae*)
497 and *Lactobacillus acidophilus* on production performance of crossbred dairy cows. *Journal of Dairy Science*,
498 **82**(Suppl. 1), 128.

- 499 Kopečný J, Hodrova B. 1995. Pectinolytic enzymes of anaerobic fungi. *Letters in Applied Microbiology*, **20**(5),
500 312-316.
- 501 Krehbiel C R, Rust S R, Zhang G, Gilliland S E. 2003. Bacterial direct-fed microbials in ruminant diets:
502 Performance response and mode of action. *Journal of Animal Science*, **81**, 120-132.
- 503 Kumar S, Choudhury P K, Carro M D, Griffith G W, Dagar S S, Puniya M, Calabro S, Ravella S R, Dhewa T,
504 Upadhyay R C, Sirohi S K, Kundu S S, Wanapat M, Puniya A K. 2014. New aspects and strategies for
505 methane mitigation from ruminants. *Applied Microbiology and Biotechnology*, **98**, 31-44.
- 506 Kumar S, Dagar S S, Puniya AK, Upadhyay R C. 2013a. Changes in methane emission, rumen fermentation in
507 response to diet and microbial interactions. *Research in Veterinary Science*, **94**, 263-268.
- 508 Kumar S, Dagar S S, Sirohi, S K, Upadhyay R C, Puniya A K. 2013b. Microbial profiles, *in vitro* gas production,
509 dry matter digestibility based on various ratio of roughage to concentrate. *Annals of Microbiology*, **63**, 541-
510 545.
- 511 Kumar S, Puniya A K, Puniya M, Dagar SS, Sirohi S K, Singh K, Griffith G W. 2009. Factors affecting rumen
512 methanogens and methane mitigation strategies. *World Journal of Microbiology and Biotechnology*, **25**, 1557-
513 1566.
- 514 Lila Z A, Mohammed N, Yasui T, Kurokawa Y, Kanda S, Itabashi H. 2004. Effects of twin strain of
515 *Saccharomyces cerevisiae* live cells on mixed ruminal microorganism fermentation *in vitro*. *Journal of*
516 *Animal Science*, **82**, 1847-1854.
- 517 Lopez S, Valdes C, Newbold C J, Wallace R J. 1999. Influence of sodium fumarate on rumen fermentation *in*
518 *vitro*. *British Journal of Nutrition*, **81**, 59-64.
- 519 Manikumar B, Puniya A K, Singh K, Sehgal J P. 2004. *In vitro* degradation of cell wall and digestibility of
520 cereal straws treated with anaerobic ruminal fungi. *Indian Journal of Experimental Biology*, **42**, 636-638.
- 521 Mann S O, Grant C, Hobson P N. 1980. Interactions of *E. coli* and lactobacilli in gnotobiotic lambs. *Microbiology*
522 *Letters*, **15**, 141-144.
- 523 Martin S A, Nisbet D J. 1992. Effect of direct-fed microbials on rumen microbial fermentation. *Journal of Dairy*
524 *Science*, **75**, 1736-1744.
- 525 McAllister T A, Bae H D, Yanke L J, Cheng K J, Muir A. 1994. Effect of condensed tannins from birds foot
526 trefoil on endoglucanase activity and the digestion of cellulose filter paper by ruminal fungi. *Canadian*
527 *Journal of Microbiology*, **40**, 298-305.
- 528 Miller-Webster T, Hoover W H, Holt M, Nocek J E. 2009. Influence of yeast culture on ruminal microbial
529 metabolism in continuous culture. *Journal of Dairy Science*, **85**, 2014- 2021.
- 530 Ishtiyak MA, Sehgal J P, Sirohi S K. 2013. Isolation and hydrolytic enzymes production potential of fungal
531 isolates from Murrah Buffaloes. *Indian Journal of Animal Nutrition*, **30**, 162-168.
- 532 Mountfort D O, Asher R A. 1989. Production of xylanase by the ruminal anaerobic fungus *Neocallimastix*
533 *frontalis*. *Applied and Environmental Microbiology* **55**(4), 1016-1022.

- 534 Mwenya B, Sar C, Pen B, Morikawa R, Takaura K, Kogawa S, Kimura K, Umetsu K, Takahashi J. 2006. Effect of
535 feed additives on ruminal methanogenesis and anaerobic fermentation of manure in cows and steers.
536 *International Congress Series*, **1293**, 209-212.
- 537 Newbold C J, Wallace R J, McIntosh F M. 1996. Mode of action of the yeast *Saccharomyces cerevisiae* as a feed
538 additive for ruminants. *British Journal of Nutrition*, **76**, 249.
- 539 Newman K E, Jacques K A. 1995. Microbial feed additives for pre-ruminants. In: *Biotechnology in Animal Feeds*
540 *and Animal Feeding*. Wallace R J, Chesson A. (Eds.), VCH, Weinheim, Germany, pp. 247–258
- 541 Nisbet D J, Martin S A. 1991. Effect of a *Saccharomyces cerevisiae* culture on lactate utilization by the ruminal
542 bacterium *Selenomonas ruminantium*. *Journal of Animal Science*, **69**, 4628.
- 543 Novotna Z, Fliegerova K, Simunek J. 2008. Characterization of chitinases of polycentric anaerobic rumen fungi.
544 *Folia Microbiologica* (Praha), **53(3)**, 241-245.
- 545 Novotna Z, Prochazka J, Simunek J, Fliegerova K. 2010. Xylanases of anaerobic fungus *Anaeromyces*
546 *mucronatus*. *Folia Microbiologica* (Praha), **55(4)**, 363-367.
- 547 Oetzel G R, Emery K M, Kautz W P, Nocek J E. 2007. Direct-fed microbial supplementation and health and
548 performance of pre- and postpartum dairy cattle: A field trial. *Journal of Dairy Science*, **90**, 2058–2068.
- 549 Ondarza de M B, Sniffen C J, Graham H, Wilcock P. 2010. Case study: Effect of supplemental live yeast on yield
550 of milk and milk components in high-producing multiparous Holstein cows. *Professional Animal Scientist*, **26**,
551 443-449.
- 552 Paul S S, Kamra D N, Sastry V R., Sahu N P, Agarwal N. 2004. Effect of anaerobic fungi on *in vitro* feed
553 digestion by mixed rumen microflora of buffalo. *Reproduction Nutrition and Development*, **44**, 313-319.
- 554 Paul S S, Kamra D N, Sastry V R., Sahu N P. 2006. Effect of adding an anaerobic fungal culture isolated from a
555 wild blue bull (*Boselophus tragocamelus*) to rumen fluid from buffaloes on *in vitro* fibrolytic enzyme activity,
556 fermentation and degradation of tannins and tannin-containing Kachnar tree (*Bauhinia variegata*) leaves and
557 wheat straw. *Journal of Science and Food Agriculture*, **86**, 258-270.
- 558 Plata F P, Mendoza G D, Barcena-Gama J R, Gonzalez S M. 1994. Effect of a yeast culture (*Saccharomyces*
559 *cerevisiae*) on neutral detergent fiber digestion in steers fed oat straw based diets. *Animal Feed Science and*
560 *Technology*, **49**, 203-210.
- 561 Qiu, R, Croom, J, Ali, R A, Ballou, A L, Smith, C D, Ashwell C M, Hassan, H M, Chiang C-C, Koci, M D.
562 (2012). Direct fed microbial supplementation repartitions host energy to the immune system. *Journal of*
563 *Animal Science*, **90**, 2639-2651.
- 564 Raeth-Knight, M L, Linn, J G, Jung H G. (2007).Effect of Direct-Fed Microbials on Performance, Diet
565 Digestibility, and Rumen Characteristics of Holstein Dairy Cows. *J. Dairy Science*, **90**, 1802-1809.
- 566 Riddell J B, Gallego A J, Harmon D L, McLeod K R. 2010. Addition of a *Bacillus* based probiotic to the diet of
567 preruminant calves: Influence on growth, health, and blood parameters. *International Journal of Applied*
568 *Research in Veterinary Medicine*, **8**, 78-85.
- 569 Sadine W E. 1979. Roles of lactobacillus in the intestinal tract. *Journal of Food Production*, **42**, 259-262.

- 570 Sakurada M, Morgavi D P, Tomita Y, Onodera R. 1995. Chitinolytic activity of the anaerobic rumen fungus
571 *Piromyces communis*. *Current Microbiology* **31**(4), 206-209.
- 572 Sandri M, Manfrin C, Pallavicini A, Stefanon B. 2014. Microbial diversity of the liquid fraction of rumen content
573 from lactating cows. *Animal*, 1-8.
- 574 Saxena S, Sehgal J, Puniya A K, Singh K. 2010. Effect of administration of rumen fungi on production
575 performance of lactating buffaloes. *Beneficial Microbes*, **1**, 183-188.
- 576 Schingoethe D J, Linke K N, Kalscheur K F, Hippen A R. 2004. Feed efficiency of mid-lactation dairy cows fed
577 yeast culture during summer. *Journal of Dairy Science*, **87**, 4178–4181.
- 578 Sehgal J P, Jit D, Puniya A K, Singh K. 2008. Influence of anaerobic fungal administration on growth, rumen
579 fermentation and nutrient digestion in female buffalo calves. *Journal of Animal Feed Science*, **17**, 510–518.
- 580 Shelke S K, Chhabra A, Puniya A K, Sehgal J P. 2009. *In vitro* degradation of sugarcane bagasse based ruminant
581 rations using anaerobic fungi. *Annals of Microbiology*, **59**(3), 415-418.
- 582 Teather R M, Froster R J. 1998. Manipulating the rumen microflora with bacteriocin to improve ruminant
583 production. *Canadian Journal of Animal Science*, **78**, 57–69.
- 584 Teunissen MJ, Op den Camp HJ. 1993. Anaerobic fungi and their cellulolytic and xylanolytic enzymes. *Antonie*
585 *Van Leeuwenhoek*, **63**(1), 63-76.
- 586 Thareja A, Puniya A K, Goel G, Nagpal R, Sehgal J P, Singh P, Singh K. 2006. *In vitro* degradation of wheat
587 straw by anaerobic fungi from small ruminants. *Archives of Animal Nutrition*, **60**, 412-417.
- 588 Theodorou M K, Beever D E, Haines M J, Brooks A. 1990. The effect of a fungal probiotic on intake and
589 performance of early weaned calves. *Animal Production*, **50**, 577 (Abstr).
- 590 Timmerman H M, Mulder L, Everts H, van Espen D C, van der Wal E, Klaassen G, Rouwers S M G, Hartemink
591 R, Rombouts F M, Beynen A C. 2005. Health and growth of veal calves fed milk replacers with or without
592 probiotics. *Journal of Dairy Science*, **88**, 2154- 2165.
- 593 Trinci A P J, Davies D R, Gull K, Lawrence M I, Nielsen B B, Rickers A, Theodorou M K. 1994. Anaerobic fungi
594 in herbivorous animals. *Mycological Research*, **98**, 129–152.
- 595 Tripathi V K, Sehgal J P, Puniya A K, Singh K. 2007a. Hydrolytic activities of anaerobic fungi isolated from wild
596 blue bull (*Boselaphus tragocamelus*). *Anaerobe*, **13**, 36-39.
- 597 Tripathi V K, Sehgal J P, Puniya A K, Singh K. 2007b. Effect of administration of anaerobic fungi isolated from
598 cattle and wild blue bull (*Boselaphus tragocamelus*) on growth rate and fibre utilization in buffalo calves.
599 *Archives of Animal Nutrition*, **61**, 416-423.
- 600 Vandevoorde L, Christianens H, Verstraete W. 1991. *In vitro* appraisal of the probiotic value of intestinal
601 lactobacilli. *World Journal of Microbiology and Biotechnology*, **7**, 587-592.
- 602 Varel V H, Kreikemeier K K, Jung H J G, Hatfield R D. 1993. *In vitro* stimulation of forage fiber degradation by
603 ruminal microorganisms with *Aspergillus oryzae* fermentation extract. *Applied and Environmental*
604 *Microbiology*, **59**, 3171-3176.

- 605 Waldrip H M, Martin S A. 1993. Effects of an *Aspergillus oryzae* fermentation extract and other factors on lactate
606 utilization by the ruminal bacterium *Megasphaera elsdenii*. *Journal of Animal Science*, **71**, 2770-2776.
- 607 Wallace R J, Joblin K N. 1985. Proteolytic activity of a rumen anaerobic fungus. *FEMS Microbiology Letters*,
608 **29(1-2)**, 19-25.
- 609 Wallace R J, Newbold, C J. 1993. Rumen fermentation and its manipulation: The development of yeast culture as
610 feed additives. In: Lyons T P (Ed.), *Biotechnology in the Feed Industry* Kentucky, Alltech Technical
611 Publications, pp. 173-192.
- 612 Walsh M C, Rostagno M H, Gardiner G E, Sutton A L, Richert B T, Radcliff J S. 2012. Controlling *Salmonella*
613 infection in weaning pigs through water delivery of direct-fed microbials or organic acids. Part I: effect on
614 growth performance, microbial populations, and immune status. *Journal of Animal Science*, **90**, 261-271.
- 615 Wisener L V, Sargeant J M, Connor A M O, Faires M C, Glass-Kasstra S K. 2014. The Use of Direct-Fed
616 Microbials to Reduce Shedding of *Escherichia coli* O157 in Beef Cattle: A Systematic Review and Meta-
617 analysis. *Zoonosis and Public Health* (In Print)
- 618 Wohlt J E, Finkelstein A D, Chung C H. 1991. Yeast culture to improve intake, nutrient digestibility, and
619 performance by dairy cattle during early lactation. *Journal of Dairy Science*, **74**, 1395–1400.
- 620 Yoon I K, Stern M D. 1995. Influence of direct-fed microbials on ruminal microbial fermentation and
621 performance of ruminants: A review. *Asian-Australasian Journal of Animal Science*, **8**, 533–555.
- 622 Yu P, Huber J T, Theurer C B, Chen K H, Nussio L G, Wu Z. 1997. Effect of steam-flaked or steam-rolled corn
623 with or without *Aspergillus oryzae* in the diet on performance of dairy cows fed during hot weather. *Journal*
624 *of Dairy Science*, **80**, 3293–3297.
- 625 Yue Q, Yang H J, Cao Y C, Zhang D F, Jiang Y H, Wang J Q. 2009. Feruloyl and acetyl esterase production of an
626 anaerobic rumen fungus *Neocallimastix* sp. YQ2 effected by glucose and soluble nitrogen supplementations
627 and its potential in the hydrolysis of fibrous feedstuffs. *Animal Feed Science and Technology*, **153(3-4)**, 263-
628 277.
- 629

Table 1: Enzymes produced by anaerobic rumen fungi and their functions

Enzymes	Types	Function(s)	Reference(s)
Esterases	p-Coumaroyl esterase	Cleave phenolic acid (p-coumaric and ferulic acid) residues from the lignin hemicellulose or lignin xylan complexes, loosening cell wall structures, thereby allowing access to previously protected polysaccharides	Atsushi <i>et al.</i> (1984); Yue <i>et al.</i> (2009)
	Feruloyl esterase		
	Acetyl esterase	acetyl xylan esterases remove acetyl group more specifically from xylose moieties in the xylan main chain	Blum <i>et al.</i> (1999)
Cellulases	Endoglucanases	These act in synergy to convert cellulose to glucose. Initial attack on the cellulose molecule is by the endo-glucanase, which cuts the linear cellulose chains internally. Exo-glucanase can then act at these nick sites, releasing cellobiose, which is in turn hydrolysed by β -glucosidase to glucose monomers	Teunissen and Op den Camp (1993); Gordon and Phillips (1998); Atanasova-Pancevska and Kungulovski(2008); Comlekcioglu <i>et al.</i> (2010)
	Exoglucanase		
	β -glucosidase		
Hemicellulases	Xylanase	Degrade Xylan	Mountfort and Asher (1989); Teunissen and Op den Camp (1993); Breton <i>et al.</i> (1995); Blum <i>et al.</i> (1999); Novotna <i>et al.</i> (2010)
	Mannase	Degrade manose	Coughlan and Hazlewood (1993)
Pectinases	Endocellular pectin lyase		Kopečný and Hodrova (1995)
	Polygalacturonase		
Proteases		The contribution made by protease of anaerobic fungi in degradation of dietary proteins remains unclear	Wallace and Joblin (1985)
Chitinases			Sakurada <i>et al.</i> (1995); Novotna <i>et al.</i> (2008)